



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07D 209/14, A61K 31/40</b>	<b>A1</b>	(11) International Publication Number: <b>WO 99/51575</b> (43) International Publication Date: 14 October 1999 (14.10.99)
---	-----------	--

(21) International Application Number: PCT/US99/07621

(22) International Filing Date: 7 April 1999 (07.04.99)


(30) Priority Data:  
09/057,252 8 April 1998 (08.04.98) US

(71) Applicant: AMERICAN HOME PRODUCTS CORPORATION [US/US]; Five Giralda Farms, Madison, NJ 07940-0874 (US).

(72) Inventors: MEWSHAW, Richard, Eric; 21 Boxwood Drive, Princeton, NJ 08512 (US). ZHOU, Dahui; 807B Donaldson Street, Highland Park, NJ 08904 (US).

(74) Agents: NAGY, Michael, R.; American Home Products Corporation, Patent Law Dept. - 2B, One Campus Drive, Parsippany, NJ 07054 (US) et al.

(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published***With international search report.**Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.*(54) Title: N-ARYLOXYETHYL-INDOLY-ALKYLAMINES FOR THE TREATMENT OF DEPRESSION (5-HT<sub>1A</sub> RECEPTOR ACTIVE AGENTS)

## (57) Abstract

Compounds useful for alleviating symptoms of depression (5-HT<sub>1A</sub> receptor active agents) are provided which have formula (I), wherein: R<sub>1</sub> is hydrogen, lower alkyl or aryl; R<sub>2</sub> is hydrogen, lower alkyl, phenyl or substituted phenyl; X and Y are each, independently, hydrogen, lower alkyl, lower alkoxy, or halogen, or together combined with the carbon atoms to which they are attached to complete a cyclopentyl, cyclohexyl, phenyl, pyrrolyl, pyranal, pyridinyl, dihydrofuranyl, furanyl, dioxanyl, oxazolyl or isoxazolyl group; Z is hydrogen, halogen or lower alkoxy; with the proviso that when X, Y or Z represent lower alkoxy, they are not present at the ortho position; W is hydrogen, halogen, lower alkyl, cyano or a trifluoromethyl group; and n is 2-5; or pharmaceutically acceptable salts thereof.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

N-ARYLOXYETHYL-INDOLY-ALKYLAMINES FOR THE TREATMENT OF DEPRESSION. (5-HT<sub>1A</sub> RECEPTOR ACTIVE AGENTS)

5 **FIELD OF INVENTION**

The present invention relates to compounds useful for the treatment of diseases affected by disorders of the serotonin-affected neurological systems. More specifically, the present invention is directed to aryloxyethyl-indoly-alkylamine derivatives useful for the treatment of such diseases.

10

**BACKGROUND OF INVENTION**


Pharmaceuticals which enhance neurotransmission of serotonin (5-HT) are useful for the treatment of many psychiatric disorders, including depression and anxiety. The first generation of non-selective serotonin-affecting compounds operated through a variety of physiological means which caused them to possess numerous  
15 undesired side effects, such as dry mouth, blurred vision, and sedation due to multiple receptor activities. The more recently introduced compounds, i.e., the selective serotonin reuptake inhibitors (SSRIs), act predominately by inhibiting 5-HT, which is released at the synapses, from being actively removed from the synaptic cleft via a  
20 presynaptic serotonin transport carrier. As SSRIs require several weeks before they exert their full therapeutic effect, this 5-HT blockade mechanism cannot fully account for their therapeutic activity. It is speculated that this two week induction which occurs before a full antidepressant effect is observed, is due to the involvement of the 5-HT<sub>1A</sub> autoreceptors which suppress the firing activity of the 5-HT neurons, causing a  
25 dampening of the therapeutic effect. Studies suggest that after several weeks of SSRI administration, a desensitization of the 5-HT autoreceptors occurs allowing a full antidepressant effect in most patients, see Le Poul et al., Arch. Pharmacol., 352:141 (1995). Hence, it is believed that overriding this negative feedback by using 5HT<sub>1A</sub> antagonists would increase and accelerate the clinical antidepressant response. Recent  
30 studies by Artigas et al., Trends Neurosci., 19:378-383, (1996) suggest that a combination of 5-HT<sub>1A</sub> activity and inhibition of 5-HT uptake within a single molecular entity can achieve a more robust and fast-acting antidepressant effect.

The present invention relates to a new class of molecules which have the ability  
35 to act concomitantly at the 5-HT<sub>1A</sub> autoreceptors and with the 5-HT transporter. Such compounds are therefore potentially useful for the treatment of anxiety or depression, as well as other serotonin disorders.


- 2 -

U.S. Patent No. 3,371,098 discloses sec. and tert. indolyethylamines useful as sedatives, anticonvulsants and analgesics.


- 5 U.S. Patent No. 5,436,264 discloses N-aryloxyalkyl-tryptamine-like compounds of the following formula as alpha-1-adrenergic receptor antagonists for the treatment of cardiovascular disorders.



- 10 EP 0722 941 A2 discloses the preparation of a series of hetero-oxy alkanamines of the following formula for the treatment of depression and other disorders for which serotonin uptake inhibitors are normally used.




- 15 Japanese Patents 05255302 and 09040648 disclose the following compounds which are reported to be useful for the treatment of central nervous system-related diseases, such as anxiety and depression.



- 3 -

**SUMMARY OF INVENTION**

The compounds of the present invention are aminomethyl benzoxezine indoles represented by Formula I:



I

5

wherein:

R<sub>1</sub> is hydrogen, lower alkyl, or aryl;

R<sub>2</sub> is hydrogen, lower alkyl, phenyl, or substituted phenyl;

10 X and Y are each, independently, hydrogen, lower alkyl, lower alkoxy, or halogen, or together combine with the carbon atoms to which they are attached to complete a cyclopentyl, cyclohexyl, phenyl, pyrrolyl, pyranal, pyridinyl, dihydrofuranyl, furanyl, dioxanyl, oxazolyl, or isoxazolyl group;

Z is hydrogen, halogen, or lower alkoxy; with the proviso that when X, Y or Z represent lower alkoxy, they are not present at the ortho position;

15 W is hydrogen, halogen, lower alkoxy, lower alkyl, cyano, or a trifluoromethyl group; and

n is 2-5; or

pharmaceutically acceptable salts thereof.

20

The present invention is further derived to pharmaceutical compounds containing such compounds, as well as methods for alleviating symptoms of depression comprising administering the present compounds to a patient in need thereof.

**DETAILED DESCRIPTION OF THE INVENTION**

25

Preferably, the compounds of the present invention are those represented by Formula I, wherein:

R<sub>1</sub> is hydrogen, methyl or aryl;

R<sub>2</sub> is hydrogen;

30

X and Y are each, independently, hydrogen, halogen or lower alkoxy, or together combine with the carbon atoms to which they are attached to complete a cyclopentyl, cyclohexyl, phenyl, pyridinyl, dioxanyl, oxazolyl, furanyl or dihydrofuranyl group;

- 4 -

Z is hydrogen, halogen or lower alkoxy; with the proviso that when X, Y or Z are lower alkoxy they are not present at the ortho position;

W is hydrogen or halogen; and

n is 2-4; or

5 pharmaceutically acceptable salts thereof.

Most preferably, the compounds of the present invention are selected from the following:

- [3-(5-Fluoro-1H-indol-3-yl)-propyl]-[2-(1H-indol-4-yloxy)-ethyl]-amine;
- 10 [2-(1H-Indol-4-yloxy)ethyl]-[3-(1H-indol-3-yl)-propyl]-amine;
- [3-(1H-Indol-3-yl)-butyl]-[2-(1H-indol-4-yloxy)-ethyl]-amine;
- [2-(2,3-Dihydro-benzo[1,4]dioxin-5-yloxy)-ethyl]-[2-(1H-indol-3-yl)-ethyl]-amine;
- [2-(2,3-Dihydro-benzo[1,4]dioxin-5-yloxy)-ethyl]-[3-(5-fluoro-1H-indol-3-yl)-
- propyl]-amine;
- 15 [2-(6-Fluorochroman-8-yloxy)-ethyl]-[2-(1H-indol-3-yl)-ethyl]-amine;
- [2-(6-Fluorochroman-8-yloxy)-ethyl]-[3-(5-fluoro-1H-indol-3-yl)-propyl]-amine;
- [2-(6-Fluorochroman-8-yloxy)-ethyl]-[2-(5-fluoro-1H-indol-3-yl)-ethyl]-amine;
- [2-(2,3-Dihydro-benzofuran-7-yloxy)-ethyl]-[3-(5-fluoro-1H-indol-3-yl)-propyl]-amine;
- [2-(Benzofuran-7-yloxy)-ethyl]-[3-(5-fluoro-1H-indol-3-yl)-propyl]-amine;
- 20 [2-(5-Fluoro-2,3-dihydro-7-yloxy)-ethyl]-[2-(5-fluoro-1H-indol-3-yl)-ethyl]-amine;
- [3-(5-Fluoro-1H-indol-3-yl)-propyl]-[2-(indan-4-yloxy)-ethyl]-amine;
- [3-(5-Fluoro-1H-indol-3-yl)-propyl]-[2-(5,6,7,8-tetrahydro-naphthalen-1-yloxy)-
- ethyl]-amine;
- [3-(1H-Indol-3-yl)-propyl]-[2-(naphthalen-1-yloxy)-ethyl]amine;
- 25 [3-(1H-Indol-3yl)-propyl]-(2-phenoxy-ethyl)-amine;
- [3-(5-Fluoro-1H-indol-3yloxy)-propyl]-[2-(indan-5-yloxy)-ethyl]-amine;
- [3-(1H-Indol-3-yl)-propyl]-[2-(quinolin-8-yloxy)-ethyl]-amine;
- [3-(5-Fluoro-1H-indol-3-yl)-propyl]-[2-(2-methoxy-phenoxy)-2-phenyl-ethyl]-amine;
- and [3-(5-Fluoro-1H-indol-3-yl)-propyl]-[2-(2-methoxy-phenoxy)-propyl]amine;

30

As used herein, the terms "lower alkyl" and "lower alkoxy" are meant to include both straight and branched carbon chains containing 1 to 6 carbon atoms. The term "halogen" is meant to include fluorine, chlorine, bromine, and iodine. The term "substituted phenyl" is meant to include a phenyl moiety substituted with an alkyl,


35 halogen, or alkoxy group. The term "aryl" is meant to include aromatic radicals containing 6-12 carbon atoms.

- 5 -

The compounds of Formula I may advantageously be used in the form of the pharmaceutically acceptable acid addition salts thereof. Such salts, which may be prepared by methods well known to those skilled in the art, may be formed with both inorganic or organic acids, for example: fumaric, maleic, benzoic, ascorbic, pamoic, succinic, bismethylenesalicylic, methanesulfonic, ethanedisulfonic, acetic, oxalic, propionic, tartaric, salicylic, citric, gluconic, lactic, malic, mandelic, cinnamic, citraconic, aspartic, stearic, palmitic, itaconic, glycolic, p-aminobenzoic, glutamic, benzene-sulfonic, hydrochloric hydrobromic, sulfuric, cyclohexylsulfamic, phosphoric and nitric acids.


10

The compounds of the present invention may be prepared by any suitable method known to those skilled in the art. However, the present compounds may be advantageously prepared according to any one of Schemes 1 to 10 set forth below. In the Schemes, the intermediate compounds discussed hereinafter are identified in parenthesis. The compound produced in each of Schemes 1 to 10 is identified with reference to the appropriate, corresponding Example.

**Scheme 1****Ex. 1**

- 6 -


Scheme 2






- 7 -


## Scheme 3




## Scheme 4




## Scheme 5






- 10 -

Scheme 7




Scheme 8




- 12 -

## Scheme 9



## Scheme 10



- 13 -

The present invention will now be illustrated by reference to the following specific, non-limiting examples.

## INTERMEDIATE 2

### 5                    [2-(2-Methoxy-phenoxy-ethyl)-benzyl-carbamic acid    *tert*-butyl ester

To a solution of (2-hydroxy-ethyl)-benzyl-carbamic acid *tert*-butyl ester ( 3.2 g, 12.6 mmol) and 2-methoxy phenol (1.0 g, 8.4 mmol) containing triphenylphosphine (3.3 g, 12.6 mmol) in anhydrous tetrahydrofuran (40 ml) was slowly added  
10 diisopropyl azodicarboxylate (2.5 g, 12.6 mmol). The reaction was allowed to stir for 18 hours, then poured into methylene chloride (250 ml) and washed with 1N sodium hydroxide (3 x 80 ml), dried over anhydrous magnesium sulfate, filtered and concentrated to a clear oil. The oil was dissolved in ether (70 ml) and hexanes were slowly added until the triphenylphosphine oxide precipitated. The solid was filtered  
15 and the solvent removed. The thick oil was then purified by column chromatography (15% ethyl acetate-hexanes) to afford 2.57 g (57.0%) of a clear oil: MS (EI) 358 *m/e* ( $M^+$ ).

Elemental analysis for  $C_{21}H_{27}NO_4$

	Calc'd:	C, 70.56; H, 7.61; N, 3.92
20	Found:	C, 70.27; H, 7.58; N, 4.07

## INTERMEDIATE 3

### [2-(2-Methoxy-phenoxy)-ethyl]-benzyl-amine

To a solution of 2-(2-methoxy-phenoxy-ethyl)-benzyl-carbamic acid *tert*-butyl  
25 ester ( 18.0 g, 50.4 mmol) in methylene chloride (350 ml) was slowly added trifluoroacetic acid (60 ml). The reaction was stirred at room temperature for 12 hours, and then poured into 1 N sodium hydroxide (200 ml) and extracted with methylene chloride (3 x 150 ml). The combined organic layers were washed with 1 N sodium hydroxide (2 x 150 ml) followed by water (2 x 100 mL), then dried over anhydrous  
30 magnesium sulfate, filtered, and the solvent removed under vacuum. Chromatography (5% methanol-methylene chloride) afforded 12.4 g (96 %) of a clear oil.

Elemental analysis for  $C_{16}H_{19}NO_2$

	Calculated:	C, 74.19; H, 7.24; N, 5.54
	Found:	C, 73.91; H, 7.28; N, 5.44

- 14 -

The fumarate salt was prepared in ethanol: mp 121.5 -122°C:

Elemental analysis for  $C_{16}H_{19}NO_2 \cdot C_4H_4O_4$

Calc'd : C, 63.56; H, 6.27; N, 3.71

Found: C, 63.35; H, 6.16; N, 3.62

5

#### INTERMEDIATE 4

##### **N-Benzyl-3-(1H-indol-3-yl)-N-[2-(2-methoxy-phenoxy)-ethyl]-propion-amide (4a)**

To a solution of 3-indole propionic acid (4.1 g, 21.7 mmol) and 1-(3-dimethyl-  
10 amino-propyl)-3-ethylcarbodiimide hydrochloride (4.4 g, 23 mmol) in methylene  
chloride (80 ml) was added a solution of 2-(2-Methoxy-phenoxy)-ethyl]-benzyl-amine  
(3 g, 11.6 mmol) in methylene chloride (20 ml) at 0 °C. After 2 hours, the reaction  
mixture was poured into water (200 ml) and extracted with methylene chloride (2 x 50  
ml). The combined organic layers were washed with 1N sodium hydroxide (50 ml),  
15 followed by water (2 x 50 ml). The organic layer was dried over anhydrous sodium  
sulfate, filtered, and the solvent removed under vacuum. Chromatography (5%  
methanol-methylene chloride) provided 3.3 g (66.0%) of product as a white solid: mp  
46.5-47.5°C. MS EI *m/e* 428 ( $M^+$ )

20

##### **N-Benzyl-3-(1H-indol-3-yl)-N-[2-(2-methoxy-phenoxy)-ethyl]-butyr-amide (4b)**

Replacing 3-indole propionic acid with 3-indole butyric acid (1.8 g, 8.9 mmol)  
in (4a) above afforded N-benzyl-3-(1H-indol-3-yl)-N-[2-(2-methoxy-phenoxy)-ethyl]  
buty-amide (1.3 g; 78 %) as a white foam. MS FAB *m/e* 443 ( $M + H^+$ )  
25 MS FAB *m/e* 465 ( $M + Na^+$ )

#### INTERMEDIATE 5

##### **Benzyl-[3-(1H-indol-3-yl)-propyl]-[2-(2-methoxy-phenoxy)-ethyl]amine (5a)**

To a solution of N-benzyl-3-(1H-indol-3-yl)-N-[2-(2-methoxy-phenoxy)-  
ethyl]-propion-amide in tetrahydrofuran (50 mL) at room temperature was added  
lithium aluminum hydride (1.8 g). The reaction was heated to reflux for 12 hours then  
allowed to cool to room temperature. The reaction was quenched with saturated  
ammonium chloride and the solid precipitates filtered through celite. The solvent was  
35 concentrated under vacuum and the product purified by chromatography (5% methanol-  
methylene chloride) to afford 2.0 g (86%) of product as a yellow oil. MS EI *m/e* 414  
( $M^+$ )



Elemental analysis for  $C_{27}H_{30}N_2O$ 

Calc'd: C, 78.23; H, 7.30; N, 6.76

Found: C, 77.53; H, 6.95; N, 6.90

**5      Benzyl-[4-(1H-indol-3-yl)-butyl]-[2-(2-methoxyphenoxy)-ethyl]-amine (5b)**

Replacing N-benzyl-3-(1H-indol-3-yl)-N-[2-(2-methoxy-phenoxy)-ethyl]-propion-amide with N-benzyl-3-(1H-indol-3-yl)-N-[2-(2-methoxy-phenoxy)-ethyl]-butyl-amide (1.2 g, 2.7 mmol) in (5a) afforded N-benzyl-3-(1H-indol-3-yl)-N-[2-(2-methoxy-phenoxy)-ethyl]-butyl-amide (1.06 g, 91 %) as a white oil. MS EI *m/e* 428 (M<sup>+</sup>)

## INTERMEDIATE 7

### 5-Fluoro-indolyl-3-propylbromide

15 A solution of 3-(5-fluoro-1*H*-indol-3-yl)-propan-1-ol (Intermediate 6) prepared  
in accordance with the procedures set forth in Demerson et al., J. Med. Chem., 19:391-  
395 (1976). (25.4 g, 0.13 mol), carbon tetrabromide (65.5 g, 0.2 mol) and  
triphenylphosphine (52 g, 0.2 mol) in methylene chloride (156 ml) was allowed to stir  
for 2 hours. The solvent was evaporated and the product chromatographed ( 30% ethyl  
20 acetate-hexanes) to afford 33.5 g (99 %) of product.

## INTERMEDIATE 8

### 5-Fluoro-indolyl-3-propylazide

25 A solution of 5-fluoro-indolyl-3-propylbromide (10.67 g, 41 mmol) and sodium azide (3.9 g, 60 mmol) in anhydrous N,N-dimethylformamide (60 ml) was allowed to stir at 60°C for 18 hours. The mixture was poured into water (150 ml), extracted with methylene chloride (3 x 150 ml), washed with water (3 x 100 ml). The organic layer was dried over sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography (30% ethyl acetate-hexanes) afforded 8.10 g (89%) of  
30 product as a clear oil.

Elemental analysis for  $C_{11}H_{11}FN_3$ 

Calc'd: C, 60.54; H, 5.08; N, 25.67

Found: C, 60.62; H, 5.08; N, 25.84

- 16 -

**INTERMEDIATE 9****5-Fluoro-indolyl-3-propylamine**

A solution of 5-fluoro-indolyl-3-propylazide (8 g, 0.037 mol) and 10% palladium on carbon in ethanol was hydrogenated at 50 psi for 16 hours. The catalyst  
5 was filtered and the solvent removed under vacuum. The celite was washed with methanol (300 ml) and the solvent was removed under vacuum. Chromatography (15% methanol-methylene chloride plus ammonium hydroxide) afforded 4.33 g (61%) of product as a yellow solid: mp 82-84.5°C.

10 Elemental analysis for  $C_{11}H_{13}FN_2$

Calc'd: C, 68.73; H, 6.82; N, 14.57

Found: C, 68.82; H, 6.85; N, 14.49

**INTERMEDIATE 10**

15 **[3-(5-Fluoro-1H-indol-3-yl)-propyl]-[2-(2-Methoxy-phenoxy) ethyl]-benzylamine**

A solution of [2-(2-methoxy-phenoxy)-ethyl]-benzyl-amine (1.0 g, 3.9 mol), 3-(5-fluoro-1H-indol-3-yl)-propylamine (1.4 g, 5.8 mmol) and triethylamine (0.79 g, 7.8 mmol) in dimethylsulfoxide (40 ml) was allowed to stir for 16 hours at 100°C.  
20 The reaction mixture was poured into water (100 ml) and extracted with methylene chloride (3 x 100 ml). The organic layer was washed with water (3 x 100 ml), dried over anhydrous sodium sulfate, filtered, and the solvent was removed under vacuum. Chromatography (30% ethyl acetate-hexanes) afforded 0.94 g (58%) of product as a yellow oil. MS EI *m/e* 432 ( $M^+$ )

25

**INTERMEDIATE 11****2-(5-Fluoro-1H-indol-3-yl)ethanol**

To a solution of 5-fluoro-indol-3-acetic acid (4.3 g, 0.022 mol) in anhydrous THF (35 ml) was added  $LiAlH_4$  (1.0 M, 33 ml, 0.033 mol) at 0°C. The mixture was  
30 allowed to stir for 0.5 hours and, then quenched by saturated  $NH_4Cl$  solution. The mixture was then filtered through celite. The filtrate was washed with 1N NaOH (3 x 100 ml) and extracted with ethyl acetate (3 x 100 ml). The organic layer was dried over anhydrous sodium sulfate, filtered, and the solvent was removed under vacuum. Chromatography (10% methanol-methylene chloride) afforded 4.04 g (100%) of  
35 product as an off-white solid: mp 59-61°C.

Elemental analysis for  $C_{10}H_{10}FNO$

- 17 -

Calc'd: C, 67.03; H, 5.63; N, 7.82

Found: C, 66.71; H, 5.50; N, 7.74

**INTERMEDIATE 12****5 3-(5-Fluoro-1H-indol-3-yl)-ethylbromide**

To a solution of 2-(5-fluoro-1H-indol-3-yl)ethanol (4 g, 22.5 mmol) in methylene chloride (50 ml) was added carbon tetrabromide (11.2 g, 34 mmol), followed by triphenylphosphine (8.8 g, 33 mmol) at room temperature. The reaction mixture was allowed to stir at room temperature for 2.5 hours. The solvent was removed under vacuum. Chromatography (30% ethyl acetate-hexanes) afforded 5.91 g (98%) of product as an off-white solid: mp 58-59°C.

Elemental analysis for  $C_{10}H_9FBrN$ 

Calc'd: C, 49.61; H, 3.75; N, 5.79

Found: C, 49.29; H, 3.73; N, 5.72

15

**INTERMEDIATE 15****2-(5-Fluoro-2-methoxy-phenoxy)ethylchloride**

A solution of 5-fluoro-2-methoxy-phenol (4.34 g, 31 mmol) prepared in accordance with the procedures set forth in Mancini et al., Synth. Comm., 19:2001-2005 (1989), 1-bromo-2-chloroethane (8.9 ml, 107 mmol) and potassium carbonate (14.8 g, 106 mmol) in 2-butanone (60 ml) was refluxed for 24 hours. The mixture was poured into water (150 ml), extracted with methylene chloride (3 x 150 ml) and washed with brine (3 x 100 ml). The organic layer was dried over sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography (20% ethyl acetate-hexanes) afforded 4.77 g (76%) of product as a clear oil.

Elemental analysis for  $C_9H_{10}FClO_2$ 

Calc'd: C, 52.83; H, 4.93

Found: C, 52.79; H, 4.75

30

**INTERMEDIATE 16****2-(5-Fluoro-2-methoxy-phenoxy)ethylazide**

A solution of 2-(5-fluoro-2-methoxy-phenoxy)ethylchloride (3.97 g, 19 mmol) and sodium azide (2.6 g, 39 mmol) in anhydrous N,N-dimethylformide (60 ml) was allowed to stir at 60°C for 18 hours. The mixture was poured into water (150 ml), extracted with methylene chloride (3 x 150 ml), washed with water (3 x 100 ml). The organic layer was dried over sodium sulfate, filtered and the solvent was removed

35

- 18 -

under vacuum. Chromatography (20% ethyl acetate-hexanes) afforded 3.75 g (92%) of product as a clear oil.

Elemental analysis for  $C_9H_{10}FN_3O_2$

Calc'd: C, 51.18; H, 4.77; N, 19.90

5 Found: C, 51.35; H, 4.71; N, 20.06

### INTERMEDIATE 17

#### 2-(5-Fluoro-2-methoxy-phenoxy)ethylamine

A solution of 2-(5-fluoro-2-methoxy-phenoxy)ethylazide (3.97 g, 0.019 mol) and triphenylphosphine (5.95 g, 0.023 mol) in tetrahydrofuran (80 ml) and water (1.5 ml) was allowed to stir for 18 hours at room temperature. The solvent was removed under vacuum. Chromatography (ethyl acetate) removed triphenylphosphine and triphenylphosphine oxide and (25-50% methanol-ethyl acetate plus ammonium hydroxide) afforded 3.14 g (90%) of product as a clear oil. MS EI *m/e* 185 ( $M^+$ )

15

### INTERMEDIATE 18

#### 3-Indolyl-propionamide (18a)

A solution of 3-indolepropionic acid (15 g, 79 mmol), 1, 1'-carbonyldimidazole (16.7 g, 100 mmol) in anhydrous tetrahydrofuran (150 ml) was allowed to stir for 1.5 hours at room temperature. Then  $NH_3$  was bubbled through the solution for 2.5 hours at room temperature. The solvent was removed under vacuum, and the residue was dissolved in ethyl acetate (500 ml). The organic solution was washed with water (3 x 150 ml), dried over anhydrous sodium sulfate, filtered, and the solvent was removed under vacuum. The white solid was collected and dried under vacuum 10.42 g (96%); mp 124-125°C. MS EI *m/e* 188 ( $M^+$ )

25

#### 3-Indolyl-butyramide (18b)

This compound was prepared in the manner described above for (18a), using 3-indolebutyric acid and 1, 1'-carbonyldimidazole. A 96% yield was obtained as a off-white solid: mp 86-87°C.

30

### INTERMEDIATE 19

#### 3-Indolyl-propylamine (19a)

To a solution of 3-indolyl-propionamide (5 g, 24.7 mmol) in tetrahydrofuran anhydrous (150 ml) was added lithium aluminum hydride (1.0 M solution in tetrahydrofuran; 100 ml) slowly. The reaction mixture was refluxed for 3 hours, then

35

- 19 -

was quenched by adding water (4 ml), 15% sodium hydroxide (4 ml) and water (12 ml) at 0°C. The mixture was filtered through celite and concentrated under vacuum. Chromatography (10% methanol-methylene chloride plus ammonium hydroxide) afforded 4.0 g (86%) of product as a white solid: mp 58-60.5°C. MS EI *m/e* 174 ( $M^+$ )

5

### 3-Indolyl-butylamine (19b)

This compound was prepared in the manner described above for (19a) using 3-indolyl-butyramide and lithium aluminum hydride. A 75% yield was obtained as a yellow solid: mp 51-53°C.

10

## INTERMEDIATE 20

### 2-(1H-Indol-4-yloxy)ethylchloride

To a solution of 4-hydroxyindole (4 g, 30 mmol), 2-chloroethanol (4.83 g, 60 mmol), triphenylphosphine (15.7 g, 60 mmol) in anhydrous tetrahydrofuran (40 ml) was slowly added diisopropyl azodicarboxylate (12.1g, 60 mmol). The reaction was allowed to stir for 2.5 hours at room temperature, then poured into methylene chloride (250 ml), washed with water (3 x 100 ml) and dried over anhydrous sodium sulfate. This material was filtered and the solvent was removed under vacuum. Chromatography (20% hexanes-ethyl acetate) to remove triphenylphosphine, (20% methylene chloride-hexanes) afforded 2.94 g (50%) of product as a white solid: mp 69.5-72°C.

20

## INTERMEDIATE 21

### 5-Hydroxy-(2,3)-dihydrobenzo[1,4]dioxine

Pyrogallol (5 g, 0.04 mol) was dissolved in 2-butanone (600 ml) to which potassium carbonate (1.82 g, 0.013 mol) was added. The mixture was stirred at reflux while 1,2-dibromoethane (2.48 g, 1.14 ml, 0.013 mol) was slowly added dropwise. The reaction was allowed to stir overnight and then cooled to room temperature. The mixture was poured into water (100 ml) and extracted with methylene chloride (200 ml). The organic layer was dried over anhydrous sodium sulfate, filtered, and the solvent was removed under vacuum. Chromatography (5% methanol-methylene chloride) afforded 2.74 g (45%) of product as a clear oil. MS EI *m/e* 152 ( $M^+$ )

25

30

- 20 -

**INTERMEDIATE 22****5-(2-Chloroethoxy)-(2,3)-dihydrobenzo[1,4]dioxane**

To solution of 5-hydroxybenzodioxane (1.0 g, 6.5 mmol) and 2-chloroethanol (0.79 g, 9.9 mmol), triphenylphosphine (2.6 g, 9.9 mmol) in tetrahydrofuran (50 ml) was slowly added diisopropyl azidodicarbimide (DIAD) (2.0 g, 9.8 mmol). After 2 hours, another 1.5 eq of triphenylphosphine, DIAD, and 2-chloroethanol was added thereto and the mixture stirred for another 2 hours. The reaction mixture was poured into water (100 ml), and extracted with methylene chloride (100 ml). The organic layer was separated and dried over anhydrous magnesium sulfate. This material was filtered, and the solvent removed under vacuum. Chromatography (20% ethyl acetate-hexanes) afforded 1.7 g (76%) of product as a white solid: mp 70.5-72.5°C.

Elemental analysis for  $C_{10}H_{11}ClO_3$

Cal'd C, 55.96; H, 5.17

Found: C, 55.57; H, 5.20

**INTERMEDIATE 23****2-(2,3-Dihydrobenzo[1,4]dioxin-5-yloxy)ethylazide**

A solution of 5-(2-chloroethoxy)-(2,3)-dihydrobenzo[1,4]dioxane (4.6 g, 0.02 mol) and sodium azide (2.78 g, 0.043 mol) in anhydrous N,N-dimethyl-formamide (100 ml) was allowed to stir for 18 hours at 60°C. The mixture was poured into water (200 ml), extracted with methylene chloride (3 x 100 ml). The organic layer was washed with water (3 x 150 ml), dried over anhydrous sodium sulfate, filtered, and the solvent removed under vacuum. Chromatography (20% ethyl acetate-hexanes) afforded 3.43 g (72%) of product as a clear oil. MS FAB  $m/e$  221 ( $M^+$ )

**INTERMEDIATE 24****2-(2,3-Dihydrobenzo[1,4]dioxin-5-yloxy)ethylamine**

A solution of 2-(2,3-dihydrobenzo[1,4]dioxin-5-yloxy)ethylazide (3.43 g, 0.016 mol) and triphenylphosphine (6.3 g, 0.023 mol) in tetrahydrofuran (50 ml) and water (2 ml) was allowed to stir for 18 hours at room temperature. The solvent was removed under vacuum. Chromatography (30% methanol-methylene chloride plus ammonium hydroxide) afford 1.93 g (62%) of product as a yellow oil. MS FAB  $m/e$  196 ( $M + H$ )<sup>+</sup>

- 21 -

**INTERMEDIATE 25****6-Fluorochroman**

A mixture of 6-fluoro-4-chromanone (2 g, 12 mmol) and 10% palladium on carbon (1 g) in concentrated hydrochloric acid (20 ml) and ethanol (30 ml) was hydrogenated for 20 hours. The catalyst was filtered and the solvent removed under vacuum. The residue was dissolved in ethyl acetate (100 ml), washed with 1N NaOH (6 x 200 ml) and water (3 x 150 ml), dried over anhydrous sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography (20% ethyl acetate-hexanes) afforded 1.41 g (77%) of product as a clear oil. MS EI *m/e* 152 (*M*<sup>+</sup>)

10

**INTERMEDIATE 26****6-Fluorochroman-8-carbaldehyde**

To a solution of 6-fluorochroman (0.7 g, 4.6 mmol) in anhydrous methylene chloride (20 ml) was added TiCl<sub>4</sub> (1.57 g, 8.3 mmol) and  $\alpha,\alpha'$ -dichloromethyl methyl ether (0.53 g, 4.6 mmol) slowly at 0°C. The reaction was allowed to reach room temperature slowly and stirred for 16 hours. The reaction mixture was poured into ice-water, extracted with methylene chloride (3 x 100 ml), washed with saturated sodium carbonate (5 x 150 ml) and brine (3 x 100ml). The organic layer was dried over anhydrous sodium sulfate, filtered, and the solvent was removed under vacuum. The crude solid was collected and dried under the vacuum to afford 0.75 g (90%) of product as a yellow solid: mp 55-57°C.

20

Elemental analysis for C<sub>10</sub>H<sub>9</sub>FO<sub>2</sub>

Calc'd: C, 66.66; H, 5.04

Found: C, 66.64; H, 4.78

25

**INTERMEDIATE 27****6-Fluoro-8-hydroxychroman**

To a solution of 6-fluorochroman-8-carbaldehyde (8.6 g, 48 mmol), 3-*t*-butyl-4-hydroxy-5-methylphenyl sulfide (100 mg) in anhydrous methylene chloride (60 ml) at 0°C was added 3-chloroperoxybenzoic acid (mCPBA) (12.4 g, 70 mmol) portionwise. The reaction mixture was allowed to reflux for 16 hours. The excess mCPBA was destroyed by adding 10% sodium sulfite. The benzoic acid was filtered and the filtrate was extracted with methylene chloride (3 x 150 ml) and washed with water (3 x 150 ml). The organic layer was dried over anhydrous sodium sulfate and filtered. The solvent was removed under vacuum and the crude product (10.2 g, 52 mmol) was dissolved in ethanol-water (200 ml, 1:1). To the above solution, sodium

35

- 22 -

hydroxide (6.2 g, 160 mmol) was added at 0°C. After 30 minutes, the ice bath was removed, and the reaction mixture was allowed to stir for 3 hours at room temperature. Ethanol was then evaporated. The residue was neutralized with concentrated hydrochloric acid, extracted with methylene chloride (3 x 150 ml), washed with  
5 saturated sodium bicarbonate (2 x 100 ml) and brine (2 x 100 ml). The organic layer was dried over sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography (25% ethyl acetate-hexanes) afforded 6.9 g (79%) of product as a white solid: mp 62-63°C.

Elemental analysis for  $C_9H_9FO_2$

10            Calc'd:            C, 64.28; H, 5.39  
             Found:            C, 64.31; H, 5.27

### INTERMEDIATE 28

#### 2-(6-Fluorochroman-8-yloxy)ethylchloride

15            A solution of 6-fluorochroman-8-carbaldehyde (5.5 g, 33 mmol), 1-bromo-2-chloroethane (16.4 g, 114 mmol) and  $K_2CO_3$  (16 g, 114 mmol) in 2-butanone (60 ml) was refluxed for 24 hours. The mixture was poured into water (150 ml), extracted with methylene chloride (3 x 150 ml) and washed with brine (3 x 100 ml). The organic layer was dried over sodium sulfate, filtered and the solvent was removed under vacuum.  
20            Chromatography (20% ethyl acetate-hexanes) afforded 5.74 g of product as a white solid: mp 89-90°C.

Elemental analysis for  $C_{11}H_{12}FCIO_2$

             Calc'd:            C, 57.28; H, 5.24  
             Found:            C, 57.15; H, 5.69

25

### INTERMEDIATE 29

#### 2-(6-Fluorochroman-8-yloxy)ethylazide

             A solution of 2-(6-fluorochroman-8-yloxy)ethylchloride (4.13 g, 0.018 mol) and sodium azide (2.33 g, 0.036 mol) in anhydrous DMF (60 ml) was allowed to stir at  
30            60°C for 18 hours. The mixture was poured into water (150 ml), extracted with methylene chloride (3 x 150 ml) and washed with water (3 x 100 ml). The organic layer was dried over sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography (20% ethyl acetate-hexanes) afforded 4.12 g (97%) of product as a clear oil.

35            Elemental analysis for  $C_{11}H_{12}FN_3O_2$

             Calc'd:            C, 55.69; H, 5.10; N, 17.71  
             Found:            C, 55.44; H, 4.97; N, 17.88



**INTERMEDIATE 30****2-(6-Fluorochroman-8-yloxy)ethylamine**

A solution of 2-(6-fluorochroman-8-yloxy)ethylazide (4.12 g, 0.017 mol) and  
5 triphenylphosphine (6.83 g, 0.026 mol) in tetrahydrofuran (80 ml) and water (1.5 ml)  
was allowed to stir for 18 hours at room temperature. The solvent was removed under  
vacuum. Chromatography (ethyl acetate) removed triphenylphosphine and  
triphenylphosphine oxide and (40% methanol-methylene chloride plus ammonium  
hydroxide) afforded 3.45 g (94%) of product as a white solid: mp 68-70°C.

10 Elemental analysis for  $C_{11}H_{14}FNO_2$

Calc'd: C, 62.55; H, 6.68; N, 6.63

Found: C, 62.18; H, 6.54; N, 6.63

**INTERMEDIATE 31****7-Methoxybezofuran**

A solution of 7-methoxy-2-benzofurancarboxylic acid (5 g, 0.026 mol),  
copper (0.2 g) in quinoline (30 ml) was heated under reflux for 2 hours. The mixture  
was filtrated through the celite. The celite was washed with ethyl acetate. The solvent  
was removed under vacuum. Chromatography (25% ethyl acetate-hexanes) afforded  
20 2.45 g (64%) of product as a yellow oil. MS EI *m/e* 148 ( $M^+$ )

**INTERMEDIATE 32****7-Hydroxybezofuran**

7-Methoxybenzofuran (1 g, 6.7 mmol) was dissolved in anhydrous methylene  
25 chloride (25 ml) in a 100 ml round-bottom flask. The flask was placed in an acetone-  
ice bath at -78°C. The flask was fitted with a air-condenser. A solution of boron  
tribromide in methylene chloride (1 M, 10 ml) was added carefully to the stirred  
solution through the condenser. The reaction was kept at -78°C for 6 hours and then  
was allowed to stir at room temperature overnight. The reaction was quenched by  
30 adding water (20 ml) and diluted with ethyl ether. The solvent was removed under  
vacuum. Chromatography (25% ethyl acetate-hexanes) afforded 0.47 g (52%) of  
product as a light-brown oil. MS EI *m/e* 134 ( $M^+$ )

**INTERMEDIATE 33****2-(Benzofuran-7-yloxy)-ethylchloride**

35 To a solution of 7-hydroxybenzofuran (0.47 g, 3.5 mmol), triphenylphosphine  
(2.3 g, 8.7 mmol) and 2-chloroethanol (0.7 g, 8.7 mmol) in tetrahydrofuran ((50 ml)

- 24 -

was slowly added diisopropyl azodicarboxylate (1.8 g, 8.7 mmol). The reaction was stirred at room temperature for 3 hours. THF was removed under vacuum. Chromatography (25 % ethyl acetate-hexanes) afforded 0.58 g (84%) of product as a yellow oil. MS EI *m/e* 196 ( $M^+$ )

5

**INTERMEDIATE 34****2-(2,3-Dihydrobenzofuran-7-yloxy)ethylchloride**

A solution of 2-(benzofuran-7-yloxy)-ethylchloride (0.64 g) and 10% palladium on carbon in acetic acid (20 ml) was hydrogenated under 40 psi for 20 hours. The catalyst was filtered off and the solvent removed under vacuum. Chromatography (20% ethyl acetate-hexanes) afforded 0.39 g (60%) of product as a white solid: mp 49-52°C. MS EI *m/e* 198 ( $M^+$ )

10

**INTERMEDIATE 35****2-(4-Fluorophenoxy)-acetaldehyde diethyl acetal**

15

To a suspension of NaOH (5.4 g, 0.134 mol) in anhydrous DMF (100 ml) was added 4-fluorophenol (10 g, 0.089 mol) at 0°C. After H<sub>2</sub> evolution had ceased, bromoacetaldehyde diethyl acetal (16 ml, 0.11 mol) was added. The reaction was heated at 160-170°C for 18 hours. The mixture was poured into ice-water, extracted with ethyl acetate (3 x 150 ml), washed with 1N NaOH (3 x 100 ml) and brine (3 x 100 ml). The organic layer was dried over anhydrous sodium sulfate and filtered. The solvent was removed under vacuum. Chromatography (25% ethyl acetate-hexanes) afforded 16.36 g (80%) of product as a clear oil. MS EI *m/e* 228 ( $M^+$ )

20

25

**INTERMEDIATE 36****5-Fluorobenzofuran**

To a mixture of benzene (200 ml) containing polyphosphoric acid (7.9 g, 0.035 mol) was added 2-(4-fluoro-phenoxy)-acetaldehyde diethyl acetal (8 g, 0.035 mol). The mixture was stirred vigorously while being heated to reflux for 2.5 hours. The reaction mixture was cooled to room temperature and decanted from the polyphosphoric acid. The solvent was removed under vacuum. Chromatography (5% ethyl acetate-hexanes) afforded 3.4 g (45%) of product as a clear oil.

30

**INTERMEDIATE 37****5-Fluoro-2,3-dihydrobenzofuran**

35

A solution of 5-fluorobenzofuran and 10% palladium on carbon in acetic acid (25 ml) was hydrogenated under 50 psi for 12 hours. The catalyst was filtered through

- 25 -

celite, and the celite was washed with methylene chloride (200 ml). The organic layer was washed with 1N NaOH (3 x 100 ml), brine (3 x 100 ml) and dried over anhydrous sodium sulfate. The solvent was removed under vacuum to afford 2.59 g (85%) of product as a clear oil.

5

### INTERMEDIATE 38

#### 5-Fluoro-2,3-dihydrobenzofuran-7-carbaldehyde

To a solution of 5-fluoro-2,3-dihydrobenzofuran (7 g, 0.051 mol) in anhydrous methylene chloride (40 ml) was added  $\text{TiCl}_4$  (9.5 ml, 0.087 mol), followed by  $\alpha, \alpha'$ -dichloromethyl methyl ether (4.6 ml, 0.051 mol) at 0°C. The reaction was allowed to reach room temperature slowly and stirred overnight. The reaction mixture was poured into ice-water slowly, extracted with methylene chloride (3 x 100 ml) and washed with saturated sodium carbonate (5 x 100 ml), and brine (3 x 100 ml). The organic layer was dried over anhydrous sodium sulfate and filtered. Chromatography (25% ethyl acetate-hexanes) afforded 3.29 g (39%) of product as a white solid: mp 103-104°C.

15

Elemental analysis for  $\text{C}_9\text{H}_7\text{FO}_2$

Calc'd: C, 65.06; H, 4.75

Found: C, 65.01; H, 4.03

20

### INTERMEDIATE 39

#### 5-Fluoro-7-hydroxy-2,3-dihydro-benzofuran

To a solution of 5-fluoro-2,3-dihydrobenzofuran-7-carbaldehyde (3.29 g, 20 mmol), 3-t-butyl-4-hydroxy-5-methylphenyl sulfide (100 mg) in anhydrous methylene chloride (40 ml) at 0°C was added 3-chloroperoxybenzoic acid (mCPBA) (8.5 g, 30 mmol) portionwise. The reaction mixture was refluxed for 16 hours. The excess mCPBA was destroyed by adding 10% sodium sulfite. The benzoic acid was filtered off and the filtrate was extracted with methylene chloride (3 x 100 ml) and washed with water (3 x 100 ml). The organic layer was dried over anhydrous sodium sulfate and filtered. The solvent was removed under vacuum and crude product was dissolved in ethanol-water (100 ml, 1:1). To the above solution, sodium hydroxide (2.11 g, 53 mmol) was added at 0°C. After 30 minutes, the ice bath was removed, and the reaction mixture was allowed to stir for 3 hours at room temperature. Ethanol was evaporated and the residue was neutralized with concentrated hydrochloric acid. This mixture was extracted with methylene chloride (3 x 100 ml) and washed with saturated sodium bicarbonate (2 x 100 ml) and brine (2 x 100 ml). The organic layer was dried over sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography

35

- 26 -

(30% ethyl acetate-hexanes) afforded 1.62 g (50%) of product as a white solid: mp 102.5-103.5°C.

Elemental analysis for  $C_8H_7FO_2$

Calc'd: C, 62.34; H, 4.58

5 Found: C, 62.19; H, 4.59

#### INTERMEDIATE 40

##### 2-(5-Fluoro-2,3-dihydro-benzofuran-7-yloxy)ethylchloride

A solution of 5-fluoro-7-hydroxy-2,3-dihydro-benzofuran (1.6 g, 10 mmol), 1-bromo-2-chloroethane (7.8 g, 55 mmol) and  $K_2CO_3$  (2.2 g, 16 mmol) in 2-butanone (40 ml) was refluxed for 24 hours. The mixture was poured into water (150 ml), extracted with methylene chloride (3 x 150 ml) and washed with brine (3 x 100 ml). The organic layer was dried over anhydrous sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography (25% ethyl acetate-hexanes) afforded 15 2.10 g of product as a white solid: mp 72.5-74.5°C.

Elemental analysis for  $C_{10}H_{10}FCIO_2$

Calc'd: C, 55.44; H, 4.65

Found: C, 55.37; H, 4.58

20

#### INTERMEDIATE 41

##### 2-(5-Fluoro-2,3-dihydrobenzofuran-7-yloxy)ethylazide

A solution of 2-(5-fluoro-2,3-dihydro-benzofuran-7-yloxy)ethylchloride (2.05 g, 9.4 mmol) and sodium azide (1.23 g, 19 mol) in anhydrous DMF (30 ml) was allowed to stir at 60°C for 24 hours. The mixture was poured into water (100 ml), 25 extracted with methylene chloride (3 x 150 ml), washed with water (3 x 100 ml). The organic layer was dried over sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography (20% ethyl acetate-hexanes) afforded 2.0 g (95%) of product as a clear oil. MS ESI  $m/e$  241  $[M + 1]^+$

30

#### INTERMEDIATE 42

##### 2-(5-Fluoro-2,3-dihydrofuran-7-yloxy)ethylamine

A solution of 2-(5-fluoro-2,3-dihydrobenzofuran-7-yloxy)ethylazide (1.98 g, 89 mmol) and triphenylphosphine (2.8 g, 10.6 mmol) in tetrahydrofuran (50 ml) and water (1.5 ml) was allowed to stir for 18 hours at room temperature. The solvent was removed under vacuum. Chromatography (ethyl acetate) removed triphenylphosphine and triphenylphosphine oxide and (40% methanol-methylene chloride plus ammonium hydroxide) afforded 2.0 g (100%) of product as a clear oil. MS ESI  $m/e$  198  $[M + 1]^+$

35

- 27 -

**INTERMEDIATE 44****4-Indanol**

A solution of 1 g of 2-(5-fluoro-2,3-dihydrofuran-7-yloxy)ethylamine  
5 (Intermediate 43) (Intermediate 44), prepared in the manner described by Ross et al., J. Am. Chem. Soc., 110:6471-6480 (1988) and 10% palladium on carbon in acetic acid (50 ml), was hydrogenated for 36 hours. The catalyst was filtered off through celite. The celite was washed with methylene chloride. The organic layer was washed with saturated sodium carbonate (3 x 150 ml), brine (3 x 150 ml) and dried over sodium  
10 sulfate. The solvent was removed under vacuum. Chromatography (30% ethyl acetate-hexanes) afforded 0.78 g (86%) of product as a clear oil. MS EI *m/e* 134 ( $M^+$ )

**INTERMEDIATE 45****2-(Indan-4-yloxy)-ethylchloride (45a)**

15 To a solution of 4-indanol (0.75 g, 5.6 mmol), triphenylphosphine (4.4 g, 16.8 mmol), 2-chloroethanol (0.7 g, 8.7 mmol) in tetrahydrofuran (50 ml) was slowly added diisopropyl azodicarboxylate (1.8 g, 8.7 mmol). The reaction was stirred at room temperature for 3 hours and the solvent was removed under vacuum. Chromatography (25 % ethyl acetate-hexanes) afforded 0.58 g (84%) of product as a yellow oil. MS EI  
20 *m/e* 196 ( $M^+$ )

**2-(5,6,7,8-Tetrahydronaphthalen-1-yloxy)-ethylchloride (45b)**

This compound was prepared in the manner described for 45(a) above by replacing 4-indanol with 5,6,7,8-tetrahydro-1-naphthol (4 g, 0.027 mol) ) in 46% yield  
25 (2.57 g) as a clear oil.

Elemental analysis for  $C_{12}H_{15}ClO$

Calc'd: C, 68.41; H, 7.17

Found: C, 68.37; H, 7.25

30 **2-(Naphthalen-1-yloxy)-ethylchloride (45c)**

This compound was prepared in the manner described for 45(a) above by replacing 4-indanol with 1-naphthol (5 g, 0.035 mol) ) in 82% yield (6.34 g) as a clear oil.

Elemental analysis for  $C_{12}H_{11}ClO$

35 Calc'd: C, 69.74; H, 5.37

Found: C, 69.64; H, 5.30

- 28 -

**2-Phenoxy-ethylchloride (45d)**

This compound was prepared in the manner described above for 45(a) by replacing 4-indanol with phenol (5 g, 0.053 mol) ) in 12% yield (1.03 g) as a clear oil. MS EI *m/e* 156 ( $M^+$ )

5

**INTERMEDIATE 46****2-(Indan-5-yloxy)-ethylchloride**

A solution of 5-indanol (5 g, 0.037 mol), 1-bromo-2-chloroethanol (8.02 g, 0.056 mol) and potassium carbonate (7.7 g, 0.056 mol) in 2-butanone (40 ml) was refluxed for 18 hours. The mixture was poured into water (100 ml), extracted with methylene chloride (3 x 150 ml) and washed with water (3 x 100 ml). The organic layer was dried over anhydrous sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography (20% ethyl acetate-hexanes) afforded 3.2 g (43%) of product as a white solid: mp 45-46°C.

15 Elemental analysis for  $C_{11}H_{13}ClO$ 

Calc'd: C, 67.18; H, 6.66

Found: C, 67.03; H, 6.57

**INTERMEDIATE 47**20 **[3-(1H-Indol-3-yl)-propyl]-(2-hydroxyethyl)amine**

To a solution of 3-(1H-indol-3-yl)-propyl-amine (3.5 g, 18.6 mmol), 2-chloroethanol (1 g, 12.4 mmol) in anhydrous dimethylsulfoxide (20 ml) was allowed to stir at 80°C for 12 hours. The mixture was poured into water (100 ml), extracted with methylene chloride (3 x 150 ml) and washed with water (3 x 100 ml). The organic layer was dried over anhydrous sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography (15% methanol-methylene chloride) afforded 1.04 g (38%) of product as a yellow oil. MS EI *m/e* 218 ( $M^+$ )

25

**INTERMEDIATE 48**30 **(2-Hydroxy-ethyl)-[3-(1H-indol-3-yl)-propyl]-carbamic acid-tert-butyl ester**

A solution [3-(1H-indol-3-yl)-propyl]-(2-hydroxyethyl)amine (1.05 g, 4.5 mmol), di-*tert*-dicarbonate (5 g, 24 mmol) in anhydrous tetrahydrofuran (20 ml) was heated at 80°C for 2 hours. The mixture was poured into water (100 ml), extracted with methylene chloride (3 x 150 ml) and washed with water (3 x 100 ml). The organic layer was dried over anhydrous sodium sulfate, filtered and the solvent was removed

35

- 29 -

under vacuum. Chromatography (5% methanol-methylene chloride) afforded 0.86 g (56%) of product as a yellow oil. MS EI *m/e* 318 ( $M^+$ )

#### INTERMEDIATE 49

5

##### 2-Chloro-1-phenylethanol

A solution of 2-chloroacetophenone (5 g, 0.032 mol), sodium borohydride (6.1 g, 0.16 mol) in tetrahydrofuran (30 ml) was allowed to stir at 60°C for 18 hours. The reaction was quenched by water (200 ml) and stirred for another 2 hours. The mixture was extracted with methylene chloride (3 x 100 ml) and washed with water (3 x 150 ml). The organic layer was dried over sodium sulfate and filtered. The solvent was removed under vacuum and afforded 5.07 g (100%) of product as a clear oil. MS EI *m/e* 156 ( $M^+$ )

#### INTERMEDIATE 50

15

##### 2-(2-Methoxy-phenoxy)-2-phenyl-ethylchloride

To a solution of guaiacol (2 g, 8.1 mmol), triphenylphosphine (6.4 g, 24 mmol), and 2-chloro-1-phenylethanol (3.78 g, 24 mmol) in tetrahydrofuran (50 ml) was slowly added diisopropyl azodicarboxylate (4.8 g, 24 mmol). The reaction was stirred at room temperature for 2 hours. Tetrahydrofuran was removed under vacuum. Chromatography (20 % ethyl acetate-hexanes) afforded 3.30 g (78%) of product as a clear oil. MS EI *m/e* 262 ( $M^+$ )

#### INTERMEDIATE 51

##### 5-Fluoro-3-(3-p-toluenesulfonyloxypropyl)indole

25

To a stirred solution of 3-(5-fluoro-1H-indol-3-yl)-propan-1-ol (2.90 g, 15.0 mmol) in pyridine (15.0 mL) was added p-toluenesulfonyl chloride (7.1 g, 37.5 mmol) at room temperature. After stirring for 30 minutes at room temperature, the reaction mixture was poured to 200 mL of ice water. The aqueous was extracted with ethyl acetate and the combined organic extracts were washed with 1N HCl and brine. The resulting material was dried over anhydrous sodium sulfate, filtered, and the solvent was concentrated under vacuum. The crude product was purified by silica gel column chromatography (ethyl acetate/hexane, 3/7) to give 4.1 g (79 %) of the title compound as a solid: mp 74 °C (Lit. mp 99 °C; EP 464604 A2).

- 30 -

**INTERMEDIATE 52****2-(2-Methoxy-phenoxy)propionitrile**

To a solution of guaiacol (5 g, 0.04 mol) in anhydrous N,N-dimethylformide (20 ml) was added sodium hydride (1.16 g, 0.048 mol). The mixture was allowed to stir at room temperature for 0.5 hours, followed by the addition of 2-bromopropionitrile (8.09 g, 0.06 mol). The mixture was allowed to stir for 4 hours at room temperature and quenched with water (20 ml). The mixture was extracted with methylene chloride (3 x 100 ml), washed with water (3 x 100 ml) and the organic layer was dried over anhydrous sodium sulfate and filtered. The solvent was removed under vacuum. Chromatography (20% ethyl acetate-hexanes) afforded 3.91 g (55%) of product as a clear oil. MS EI *m/e* 177 ( $M^+$ )

**INTERMEDIATE 53****2-(2-Methoxy-phenoxy)propylamine**

To a solution of 2-(2-methoxy-phenoxy)-propionitrile (3.91 g, 0.022 mol) in anhydrous ethyl ether (30 ml) was added lithium aluminum hydride (1.0 M, 44 ml, 0.044 mol). The mixture was heated at 65°C for 18 hours and the reaction was quenched with water (3 ml), 15% NaOH (3 ml) and water (9 ml). The resulting material was filtered through celite. The celite was washed with methanol (200 ml), and the solvent was removed under vacuum. Chromatography (10% methanol-methylene chloride) afforded 0.26 g (7%) of product as yellow oil. MS EI *m/e* 181 ( $M^+$ )

**EXAMPLE 1****[3-(1H-Indol-3-yl)-propyl]-[2-(2-methoxy-phenoxy)-ethyl]-amine**

A mixture of benzyl-[3-(1H-indol-3-yl)-propyl]-[2-(2-methoxy-phenoxy)-ethyl]amine (2 g, 4.7 mmol) and 5% palladium on carbon in ethanol was hydrogenated for 20 hours. The catalyst was filtered off and the solvent removed under vacuum. Chromatography (ethyl acetate-hexanes-methanol-ammonia hydroxide: 4/4/1/1) afforded 0.79 g (52 %) of product as a white solid: mp 101-102°C.

The fumarate salt was prepared in ethanol: mp 130-130.5°C.

Elemental analysis for  $C_{20}H_{24}N_2O_2 \cdot C_4H_4O_4$

Calc'd: C, 64.78; H, 6.46; N, 6.29

Found: C, 64.76; H, 6.23; N, 6.21



- 31 -

**[4-(1H-Indol-3-yl)-butyl]-[2-(2-methoxy-phenoxy)-ethyl]-amine**

Hydrogenation of benzyl-[4-(1H-indol-3-yl)-butyl]-[2-(2-methoxy-phenoxy)-ethyl]-amine afforded 0.79 g (100 %) of product as a clear oil. The oxalate salt was prepared from isopropanol: mp 167-168 °C.

- 5    Elemental analysis for  $C_{20}H_{24}N_2O_2 \cdot C_4H_4O_4$   
          Calc'd:        C, 64.47; H, 6.59; N, 6.54  
          Found:       C, 64.44; H, 6.52; N, 6.46

**EXAMPLE 2**

10    **[3-(5-Fluoro-1H-indol-3-yl)-propyl]-[2-(2-Methoxy-phenoxy)ethyl]-amine**

A mixture of [3-(5-fluoro-1H-indol-3-yl)-propyl]-[2-(2-methoxy-phenoxy)ethyl]-benzylamine (0.94 g, 2.2 mol) and 10% palladium on carbon (250 mg) in ethanol was hydrogenated for 20 hours. The catalyst was filtered off and the solvent  
15    was removed under vacuum. Chromatography (10% methanol-methylene chloride) afforded 0.63 g (85%) of product as an off-white solid: mp 125-126°C.

The oxalate salt was prepared in isopropanol: mp 146-149°C.

Elemental analysis for  $C_{20}H_{23}FN_2O_2 \cdot C_2H_2O_4 \cdot 0.5H_2O$

- Calc'd :        C, 59.85; H, 5.94; N, 6.35  
20        Found:        C, 60.13; H, 5.67; N, 6.10

**EXAMPLE 3**

**[3-(5-Fluoro-1H-indol-3-yl)propyl]-[2-(5-fluoro-2-methoxy-phenoxy)-ethyl]-amine**

- 25        A solution of 2-(5-fluoro-2-methoxy-phenoxy)ethylchloride (0.3 g, 1.5 mmol), 3-(5-fluoro-1H-indol-3-yl)propylamine (0.56 g, 2.9 mmol) in dimethylsulfoxide (20 ml) was allowed to stir for 12 h at 90°C. The reaction mixture was poured into water (100 ml) and extracted with methylene chloride (3 x 100 ml). The organic layer was washed with water (3 x 150 ml), dried over anhydrous sodium sulfate, filtered and the  
30    solvent was removed under vacuum. Chromatography (10% methanol-methylene chloride) afforded 0.39 g (77%) of product as a white solid: mp 119-122°C.

The oxalate salt was prepared in ethanol: mp 175-177°C

Elemental analysis for  $C_{20}H_{22}F_2N_2O_2 \cdot C_2H_2O_4$

- Calc'd:        C, 58.66; H, 5.57; N, 6.22  
35        Found:        C, 58.29; H, 5.25; N, 6.07

- 32 -

**[2-(5-Fluoro-1H-indol-3-yl)ethyl]-[2-(5-fluoro-2-methoxy-phenoxy)-ethyl]-amine (3b)**

A solution of 2-(5-fluoro-2-methoxy-phenoxy)ethylamine (0.51 g, 2.8 mmol), 2-(5-fluoro-1H-indol-3-yl)ethylchloride (0.44 g, 1.8 mmol) and triethylamine (0.29 g, 3 mmol) in dimethylsulfoxide (20 ml) was allowed to stir for 8 hours at 90°C. The reaction mixture was poured into water (100 ml) and extracted with methylene chloride (3 x 100 ml). The organic layer was washed with water (3 x 150 ml), dried over anhydrous sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography (5% methanol-methylene chloride) afforded 0.27 g (43%) of product as a brown oil.

The oxalate salt was prepared in ethanol: mp 185-188°C

Elemental analysis for  $C_{19}H_{20}F_2N_2O_2 \cdot C_2H_2O_4$

Calc'd: C, 57.80; H, 5.08; N, 6.42

Found: C, 57.53; H, 4.95; N, 6.36

**EXAMPLE 4**

**[2-(5-Fluoro-1H-indol-3-yl)ethyl]-[2-(5-fluoro-2-methoxy-phenoxy)-ethyl]-amine**

A solution of 2-(5-fluoro-2-methoxy-phenoxy)ethylamine (0.41 g, 2.2 mmol), 3-(2-bromoethyl)indole (0.25 g, 1.1 mmol) in dimethylsulfoxide (20 ml) was allowed to stir for 12 hours at 90°C. The reaction mixture was poured into water (100 ml) and extracted with methylene chloride (3 x 100 ml). The organic layer was washed with water (3 x 150 ml), dried over anhydrous sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography (10% methanol-methylene chloride) afforded 0.15 g (34%) of product as a brown oil.

The oxalate salt was prepared in ethanol: mp 188-189°C

Elemental analysis for  $C_{19}H_{21}FN_2O_2 \cdot C_2H_2O_4 \cdot 0.25H_2O$

Calc'd C, 59.04; H, 5.63; N, 6.62

Found: C, 59.68; H, 5.49; N, 6.56

**EXAMPLE 5**

**[3-(5-Fluoro-1H-indol-3-yl)propyl]-[2-(1H-indol-4-yloxy)-ethyl]-amine**

A solution of 2-(1H-indol-4-yloxy)ethylchloride (0.7 g, 3.6 mmol), 5-fluoro-indolyl-3-propylamine (1.0 g, 5.4 mmol) in dimethylsulfoxide (20 ml) was allowed to stir for 12 hours at 90°C. The reaction mixture was poured into water (100 ml) and extracted with methylene chloride (3 x 100 ml). The organic layer was washed with

- 33 -

water (3 x 150 ml), dried over anhydrous sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography (5%-10% methanol-methylene chloride) afforded 0.54 g (43%) of product as a white solid: mp 70-73°C.

The oxalate salt was prepared in ethanol: mp 183.5-185°C

- 5 Elemental analysis for  $C_{21}H_{22}FN_3O \cdot C_2H_2O_4$   
Calc'd C, 62.53; H, 5.48; N, 9.51  
Found: C, 62.31; H, 5.38; N, 9.35

**[2-(1H-Indol-4-yloxy)ethyl]-[3-(1H-indol-3-yl)-propyl]amine (5b)**

- 10 This compound was prepared in the manner described above for Example 5 using 2-(1H-indol-4-yloxy)ethylchloride and 3-indolyl-propylamine in 65% yield as an off-white solid: mp 109-111°C. The oxalate salt was prepared in isopropanol: mp 200.5-202°C.

Elemental analysis for  $C_{21}H_{23}N_3O \cdot C_2H_2O_4$

- 15 Calc'd: C, 65.19; H, 5.95; N, 9.92  
Found: C, 64.89; H, 6.00; N, 9.81

**[3-(1H-Indol-3-yl)butyl]-[2-(1H-indol-4-yloxy)ethyl]amine (5c)**

- 20 This compound was prepared in the manner described above for Example 5 using 2-(1H-indol-4-yloxy)ethylchloride and 3-indolyl-butylamine in 43% yield as an off-white solid: mp 70-73°C.

The oxalate salt was prepared in ethanol: mp 183.5-185°C.

Elemental analysis for  $C_{22}H_{25}N_3O \cdot C_2H_2O_4$

- 25 Calc'd: C, 62.58; H, 5.48; N, 9.52  
Found: C, 62.31; H, 5.38; N, 9.35

**EXAMPLE 6**

**[2-(2,3-Dihydrobenzo[1,4]dioxin-5-yloxy)-ethyl]-[2-(1H-indol-3-yl)-ethyl]amine**

- 30 A solution of 2-(2,3-dihydrobenzo[1,4]dioxin-5-yloxy) ethylamine (0.38 g, 1.9 mmol), 3-(2-bromoethyl)-indole (0.24 g, 1.1 mmol) and triethylamine (0.22 g, 2.2 mmol) in anhydrous dimethylsulfoxide (20 ml) was allowed to stir for 14 hours at 90°C. The mixture was poured into water (100 ml) and extracted with methylene chloride (3 x 100 ml). The organic layer was washed with water (3 x 150 ml), dried  
35 over anhydrous sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography (10% methanol-methylene choride plus ammonium hydroxide) afforded 0.19 g (52%) as a yellow oil.

- 34 -

The oxalate salt was prepared in ethanol: mp 197.5-198.5°C

Elemental analysis for  $C_{20}H_{22}N_2O_3 \cdot C_2H_2O_4$

Calc'd: C, 61.63; H, 5.64; N, 6.53

Found: C, 61.36; H, 5.46; N, 6.45s

5

#### EXAMPLE 7

##### **[2-(2,3-Dihydro-benzo[1,4]dioxin-5-yloxy)-ethyl]-[3-(5-fluoro-1H-indol-3-yl)-propyl]-amine**

A solution of 5-(2-chloroethoxy)-(2,3)-dihydrobenzo [1,4]dioxane (0.75 g, 3.5 mmol), 3-(5-fluoro-1H-indol-3-yl)-propylamine (1.0 g, 5.2 mmol) and triethylamine (0.35 g, 3.5 mmol) in anhydrous dimethylsulfoxide (20 ml) was allowed to stir for 14 hours at 100°C. The mixture was poured into water (100 ml) and extracted with methylene chloride (3 x 100 ml). The organic layer was washed with water (3 x 150 ml), dried over anhydrous sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography (5% -10% methanol-methylene chloride plus ammonium hydroxide) afforded 0.10 g (52%) of product as a yellow oil.

15

The fumarate salt was prepared in isopropanol: mp 189.5-190.5°C

Elemental analysis for  $C_{21}H_{23}FN_2O_3 \cdot 0.5C_4H_4O_4 \cdot 0.5H_2O$

Calc'd: C, 64.74; H, 5.88; N, 6.54

20

Found: C, 64.01; H, 5.95; N, 6.36

#### EXAMPLE 8

##### **[2-(6-Fluorochroman-8-yloxy)ethyl]-[2-(1H-indol-3-yl)ethyl]-amine**

A solution of 2-(6-fluorochroman-8-yloxy)ethylchloride (0.41 g, 2.2 mmol), 3-(2-bromoethyl)indole (0.25 g, 1.1 mmol) and triethylamine (0.23 g, 2.2 mmol) in anhydrous dimethylsulfoxide (20 ml) was allowed to stir for 12 hours at 90°C. The mixture was poured into water (100 ml) and extracted with methylene chloride (3 x 100 ml). The organic layer was washed with water (3 x 150 ml), dried over anhydrous sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography (10% methanol-methylene chloride plus ammonium hydroxide) afforded 0.15 g (34%) of product as a yellow oil.

25

30

The oxalate salt was prepared in ethanol: mp 213-214°C

Elemental analysis for  $C_{21}H_{23}FN_2O_2 \cdot C_2H_2O_4$

Calc'd: C, 62.11; H, 5.67; N, 6.30

35

Found: C, 62.26; H, 5.71; N, 6.19

- 35 -

**EXAMPLE 9****[2-(6-Fluorochroman-8-yloxy)ethyl]-[3-(5-fluro-1H-indol-3-yl)propyl]-amine (9a)**

A solution 2-(6-fluorochroman-8-yloxy)ethylchloride (0.25 g, 7.1 mmol), 3-(5-  
5 fluoro-1H-indol-3-yl)-propylamine (0.42 g, 2.2 mmol) and triethylamine (0.22 g, 2.2  
mmol) in anhydrous dimethylsulfoxide (20 ml) was allowed to stir for 14 hours at  
90°C. The mixture was poured into water (100 ml) and extracted with methylene  
chloride (3 x 100 ml). The organic layer was washed with water (3 x 150 ml), dried  
over anhydrous sodium sulfate, filtered and the solvent was removed under vacuum.  
10 Chromatography (10% methanol-methylene choride plus ammonium hydroxide)  
afforded 0.25 g (60%) as a white solid: mp 137-138.5°C.

The oxalate salt was prepared in isopropanol: mp 214-215°C

Elemental analysis for  $C_{22}H_{24}F_2N_2O_2 \cdot 1.5C_2H_2O_4$

	Calc'd:	C, 57.58; H, 5.22; N, 5.31
15	Found:	C, 57.75; H, 5.07; N, 5.51

**[2-(6-Fluorochroman-8-yloxy)ethyl]-[2-(5-fluro-1H-indol-3-yl)ethyl]-amine (9b)**

A solution of 2-(6-fluorochroman-8-yloxy)ethylchloride (0.26 g, 1.1 mmol), 2-  
20 (5-fluro-1H-indol-3-yl)-ethylamine (0.45 g, 2.1 mmol) and triethylamine (0.30 ml,  
2.1 mmol) in anhydrous dimethylsulfoxide (20 ml) was allowed to stir for 14 hours at  
90°C. The mixtrue was poured into water (100 ml) and extracted with methylene  
chloride (3 x 100 ml). The organic layer was washed with water (3 x 150 ml), dried  
over anhydrous sodium sulfate, filtered and the solvent was removed under vacuum.  
25 Chromatography (10% methanol-methylene choride plus ammonium hydroxide)  
afforded 0.19 g (48%) of product as a yellow oil.

The oxalate salt was prepared in ethanol: mp 201-203°C

Elemental analysis for  $C_{21}H_{22}F_2N_2O_2 \cdot 1C_2H_2O_4$

	Calc'd:	C, 59.70; H, 5.23; N, 6.05
30	Found:	C, 59.48; H, 5.08; N, 5.88

**EXAMPLE 10****[2-(2,3-Dihydrobenzofuran-7-yloxy)ethyl]-[3-(5-fluoro-1H-indol-3-yl)propyl]amine**

35 A solution of 2-(2,3-dihydrobenzofuran-7-yloxy)ethylchloride (0.38 g, 1.9  
mmol), 3-(5-fluro-1H-indol-3-yl)propylamine (0.93 g, 4.8 mmol), triethylamine

- 36 -

(0.48 g, 4.8 mmol) in anhydrous DMSO (20 ml) was allowed to stir at 90°C for 12 hours. The mixture was poured into water (100 ml) and extracted with methylene chloride (3 x 100 ml). The organic layer was washed with water (3 x 150 ml), dried over anhydrous sodium sulfate, filtered and the solvent was removed under vacuum.

5 Chromatography (10% methanol-methylene chloride plus ammonium hydroxide) afforded 0.52 g (77%) of product as a yellow oil.

The oxalate salt was prepared in ethanol: mp 158-160°C

Elemental analysis for  $C_{21}H_{23}FN_2O_2 \cdot C_2H_2O_4$

Calc'd: C, 63.82; H, 5.78; N, 5.95

10 Found: C, 63.45; H, 5.74; N, 5.76

### EXAMPLE 11

#### [2-(Benzofuran-7-yloxy)ethyl]-[3-(5-fluoro-1H-indol-3-yl)propyl]amine

A solution of 2-(benzofuran-7-yloxy)-ethylchloride (0.58 g, 2.9 mmol), 3-(5-  
15 fluoro-1H-indol-3-yl)propylamine (1.4 g, 7.4 mmol) and triethylamine (0.74 g, 7.4 mmol) in anhydrous DMSO (20 ml) was allowed to stir at 90°C for 12 hours. The mixture was poured into water (100 ml) and extracted with methylene chloride (3 x 100 ml). The organic layer was washed with water (3 x 150 ml), dried over anhydrous sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography  
20 (7% methanol-methylene chloride) afforded 0.31 g (30%) of product as a light-brown oil.

The oxalate salt was prepared in THF: mp 181-183°C

Elemental analysis for  $C_{21}H_{21}FN_2O_2 \cdot C_2H_2O_4$

Calc'd: C, 62.39; H, 5.24; N, 6.33

25 Found: C, 62.07; H, 5.24; N, 6.45

### EXAMPLE 12

#### [2-(5-Fluoro-2,3-dihydro-7-yloxy)-ethyl]-[2-(5-fluoro-1H-indol-3-yl)-ethyl]-amine

30 A solution of 2-(5-fluoro-2,3-dihydrofuran-7-yloxy) ethylamine (0.40 g, 2.1 mmol), 2-(5-fluoro-1H-indol-3-yl)-ethylamine (0.25 g, 1.0 mmol) and triethylamine (0.29 ml, 2.1 mmol) in anhydrous dimethylsulfoxide (20 ml) was allowed to stir for 14 hours at 90°C. The mixture was poured into water (100 ml) and extracted with methylene chloride (3 x 100 ml). The organic layer was washed with water (3 x 100  
35 ml), dried over anhydrous sodium sulfate, filtered and the solvent was removed under

- 37 -

vacuum. Chromatography (5% methanol-methylene chloride plus ammonium hydroxide) afforded 0.2 g (54%) of product of product as a yellow oil.

The oxalate salt was prepared in ethanol: mp 209.5-210.5°C

Elemental analysis for  $C_{20}H_{20}F_2N_2O_2 \cdot 1.5C_2H_2O_4$

5            Calc'd:            C, 55.98; H, 4.70; N, 5.68

             Found:            C, 55.53; H, 4.44; N, 5.72

### EXAMPLE 13

#### **[3-(5-Fluoro-1H-indol-3-yl)-propyl]-[2-(indan-4-yloxy)-ethyl]-amine (13a)**

10

A solution of 2-(indan-4-yloxy)-ethylchloride (0.58 g, 2.9 mmol), 3-(5-fluoro-1H-indol-3-yl)-propylamine (1.4 g, 7.4 mmol) and triethylamine (1 ml, 7.4 mmol) in anhydrous dimethylsulfoxide (20 ml) was allowed to stir for 12 hours at 90°C. The mixture was poured into water (100 ml) and extracted with methylene chloride (3 x 100

15

ml). The organic layer was washed with water (3 x 100 ml), dried over anhydrous sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography (7% methanol-methylene chloride) afforded 0.31 g (30%) of product as a brown oil.

The oxalate salt was prepared in tetrahydrofuran: mp 220-222°C

Elemental analysis for  $C_{22}H_{25}FN_2O \cdot C_2H_2O_4$

20            Calc'd:            C, 65.10; H, 6.15; N, 6.33

             Found:            C, 64.74; H, 6.14; N, 6.11

#### **[3-(5-Fluoro-1H-indol-3-yl)-propyl]-[2-(5,6,7,8-tetrahydro-naphthalen-1-yloxy)-ethyl]-amine (13b)**

25

This compound was prepared in the manner described above for Example 13 by replacing 2-(indan-4-yloxy)-ethylchloride with 2-(5,6,7,8-tetrahydronaphthalen-1-yloxy)-ethylchloride (0.6 g, 2.8 mmol) in 29% yield (0.3 g) as a yellow oil.

The fumarate salt was prepared in ethanol: mp 203-205°C

Elemental analysis for  $C_{23}H_{27}FN_2O \cdot 0.5C_4H_4O_4$

30            Calc'd:            C, 70.73; H, 6.89; N, 6.60

             Found:            C, 70.49; H, 6.87; N, 6.52

#### **[3-(1H-indol-3-yl)-propyl]-[2-(naphthalen-1-yloxy)-ethyl]-amine (13c)**

This compound was prepared in the manner described above for Example 13 by replacing 2-(indan-4-yloxy)-ethylchloride with 2-(naphthalen-1-yloxy)-ethylchloride (0.6 g, 2.8 mmol), and 3-(5-fluoro-1H-indol-3-yl)-propylamine with 3-(1H-indol-3-

35

- 38 -

yl)-propylamine (1.35 g, 7.2 mmol) in 85% yield (1.42 g) as a yellow solid: mp 119-120°C.

The fumarate salt was prepared in ethanol: mp 203-205°C

Elemental analysis for  $C_{23}H_{24}N_2O \cdot 0.5C_4H_4O_4 \cdot 0.25H_2O$

5            Calc'd:            C, 73.78; H, 6.56; N, 6.88

             Found:            C, 73.74; H, 6.47; N, 6.89

### **[3-(1H-indol-3-yl)-propyl]-[2-phenoxy-ethyl]-amine (13d)**

This compound was prepared in the manner described above for Example 13 by  
10 replacing 2-(naphthalen-1-yloxy)-ethylchloride with 2-phenoxy-ethylchloride (0.59 g, 3.8 mmol) in 100% yield (1.08 g) as a yellow oil.

The fumarate salt was prepared in ethanol: mp 203-205°C

Elemental analysis for  $C_{19}H_{22}N_2O \cdot C_4H_4O_4$

             Calc'd:            C, 67.30; H, 6.38; N, 6.82

15            Found:            C, 67.01; H, 6.30; N, 6.73

### **EXAMPLE 14**

#### **[3-(5-Fluoro-1H-indol-3-yloxy)-proyl]-[2-(indan-5-yloxy)-ethyl]-amine**

A solution of 2-(indan-5-yloxy)-ethylchloride (0.7 g, 3.5 mmol), 3-(5-fluoro-  
20 1H-indol-3-yl)-propylamine (1.01 g, 5.3 mmol) and triethylamine (0.53 g, 5 mmol) in anhydrous dimethylsulfoxide (20 ml) was allowed to stir for 12 hours at 90°C. The mixture was poured into water (100 ml) and extracted with methylene chloride (3 x 100 ml). The organic layer was washed with water (3 x 100 ml), dried over anhydrous sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography  
25 (5-10% methanol-methylene chloride) afforded 0.53 g (43%) of product as a yellow oil.

The fumarate salt was prepared in ethanol: mp 179-180°C

Elemental analysis for  $C_{22}H_{25}FN_2O \cdot 0.5C_4H_4O_4$

             Calc'd:            C, 69.46; H, 6.68; N, 6.75

             Found:            C, 69.19; H, 6.67; N, 6.72

30

### **EXAMPLE 15**

#### **[3-(1H-Indol-3-yl)-propyl]-[2-(quinolin-8-yloxy)ethyl]-amine**

To a solution of (2-hydroxy-ethyl)-[3-(1H-indol-3-yl)-propyl]- carbamic acid-  
tert-butyl ester (0.86 g, 2.7 mmol), triphenylphosphine (0.71 g, 2.7 mmol) and 8-  
35 hydroxy-quinoline (0.26 g, 2.7 mmol) in tetrahydrofuran (50 ml) was slowly added diisopropyl azodicarboxylate (0.55 g, 2.7 mmol). The reaction was stirred at room



- 39 -

temperature for 3 hours. Tetrahydrofuran was removed under vacuum. Chromatography (5 % methanol-methylene chloride) afforded a yellow solid, which was dissolved in methylene chloride (30 ml). To this solution was added a trifluoroacetic acid solution (4 ml in 10 ml methylene chloride). The reaction mixture was allowed to stir for 2 h at room temperature. The mixture was then quenched with saturated sodium carbonate and extracted with methylene chloride (3 x 100 ml). The organic layer was dried over anhydrous sodium sulfate and filtered. The solvent was removed under vacuum. Chromatography (10-15% methanol-methylene plus ammonium hydroxide) afforded 0.17 g (18%) of product as a light-yellow oil.

10 The HCl salt was prepared in ethyl acetate: mp 83-86°C

Elemental analysis for  $C_{22}H_{23}N_3O \cdot HCl \cdot 1.25H_2O$

Calc'd: C, 65.34; H, 6.60; N, 10.39

Found: C, 65.24; H, 6.69; N, 10.47

15

### EXAMPLE 16

#### **[3-(5-Fluoro-1H-indol-3-yl)-propyl]-[2-(2-methoxy-phenoxy)-2-phenyl-ethyl]-amine**

A solution of 2-(2-methoxy-phenoxy)-2-phenyl-ethylchloride (0.71 g, 2.7 mmol), 2-(5-fluoro-1H-indol-3-yl)-ethylamine (1.04 g, 5.4 mmol) and triethylamine (0.75 ml, 5.4 mmol) in anhydrous dimethylsulfoxide (20 ml) was allowed to stir for 12 hours at 90°C. The mixture was poured into water (100 ml) and extracted with methylene chloride (3 x 100 ml). The organic layer was washed with water (3 x 100 ml), dried over anhydrous sodium sulfate, filtered, and the solvent was removed under vacuum. Chromatography (7% methanol-methylene chloride plus ammonium hydroxide) afforded 0.15 g (13%) of product as a yellow oil.

The citrate salt was prepared in ethyl ether: mp 64.5-67°C

Elemental analysis for  $C_{20}H_{20}F_2N_2O_2 \cdot C_6H_8O_7 \cdot 1.5H_2O$

Calc'd: C, 60.27; H, 6.01; N, 4.39

Found: C, 59.96; H, 6.00; N, 4.39

30

### EXAMPLE 17

#### **[3-(5-Fluoro-1H-indol-3-yl)-propyl]-[2-(2-methoxy-phenoxy)propyl]amine**

A solution of 2-(2-methoxy-phenoxy)propylamine (0.26 g, 1.2 mmol), 5-fluoro-3-(3-p-toluenesulfonyloxypropyl)indole (0.26 g, 0.81 mol) and triethylamine (0.13 g, 1.2 mmol) in anhydrous dimethylsulfoxide (20 ml) was allowed to stir for 8

- 40 -

hours at 90°C. The mixture was poured into water (100 ml) and extracted with methylene chloride (3 x 100 ml). The organic layer was washed with water (3 x 100 ml), dried over anhydrous sodium sulfate, filtered and the solvent was removed under vacuum. Chromatography (10% methanol-methylene chloride plus ammonium hydroxide) afforded 0.1 g (36%) of product of product as a yellow oil.

The oxalate salt was prepared in 1-propanol: mp 143-146°C

Elemental analysis for  $C_{20}H_{20}F_2N_2O_2 \cdot C_2H_2O_4 \cdot 1 H_2O$

Calc'd: C, 59.47; H, 6.29; N, 6.03

Found: C, 59.34; H, 5.87; N, 5.96

10

The activity of the present compounds is demonstrated by the following standard pharmacological test procedures.

The PCR cloning of the human 5-HT<sub>1A</sub> receptor subtype from a human genomic library has been described previously Chanda et al., Mol. Pharmacol., 43:516 (1993). A stable Chinese hamster ovary cell line expressing the human 5-HT<sub>1A</sub> receptor subtype (5-HT<sub>1A</sub>-CHO cells) was employed throughout this study. Cells were maintained in DMEM supplemented with 10% fetal calf serum, non-essential amino acids and penicillin/ streptomycin.

20

Cells were grown to 95-100% confluency as a monolayer before membranes were harvested for binding studies. Cells were gently scraped from the culture plates, transferred to centrifuge tubes, and washed twice by centrifugation (2000 rpm for 10 min., 4°C) in buffer (50 mM Tris; pH 7.5). The resulting pellets were aliquoted and placed at -80°C. On the day of assay, the cells were thawed on ice, and resuspended in buffer. Studies were conducted using [<sup>3</sup>H]8-OH-DPAT as the radioligand. The binding assay was performed in 96 well microtiter plates in a final total volume of 250 µL of buffer. Competition experiments were performed by using 7 concentrations of unlabelled drug and a final ligand concentration of 1.5 nM. Non-specific binding was determined in the presence of 10 µM 5HT. Saturation analysis was conducted by using [<sup>3</sup>H]8-OH-DPAT at concentrations ranging from 0.3-30 nM. Following a 30 minute incubation at room temperature, the reaction was terminated by the addition of ice cold buffer and rapid filtration using a M-96 Brandel Cell Harvester (Gaithersburg, MD) through a GF/B filter presoaked for 30 minutes in 0.5% polyethyleneimine.

35

- 41 -

A protocol similar to that used by Cheetham et al., Neuropharmacol. 32:737, (1993) was used to determine the affinity of compounds for the serotonin transporter. Briefly, frontal cortical membranes prepared from male Sprague-Dawley rats were incubated with  $^3\text{H}$ -paroxetine (0.1 nM) for 60 min at 25°C. All tubes also contained  
5 either vehicle, test compound (one to eight concentrations), or a saturating concentration of fluoxetine (10  $\mu\text{M}$ ) to define specific binding. All reactions are terminated by the addition of ice cold Tris buffer followed by rapid filtration using a Tom Tech filtration device to separate bound from free  $^3\text{H}$ -paroxetine. Bound radioactivity was quantitated using a Wallac 1205 Beta Plate<sup>®</sup> counter. Nonlinear regression analysis was used to  
10 determine  $\text{IC}_{50}$  values which were converted to  $\text{K}_i$  values using the method disclosed in Cheng and Prusoff, Biochem. Pharmacol., 22:3099, (1973);  $\text{K}_i = \text{IC}_{50}/((\text{Radioligand conc.})/(1 + \text{KD}))$ .

The [ $^{35}\text{S}$ ]-GTP $\gamma\text{S}$  binding assay was similar to that disclosed by Lazareno and  
15 Birdsall, Br. J. Pharmacol., 109:1120, (1993). Briefly, 5-HT $_{1A}$  cloned receptor membrane fragments (as used for 5-HT $_{1A}$  receptor binding assays) were stored at -70 °C until needed. When needed, membranes were rapidly thawed, centrifuged at 40,000 x g for 10 minutes and resuspended at 4 °C for 10 minutes in assay buffer (25 mM HEPES, 3 mM  $\text{MgCl}_2$ , 100 mM NaCl, 1 mM EDTA, 10  $\mu\text{M}$  GDP, 500 mM DTT, pH  
20 8.0). These membranes were then incubated for 30 min at 30 °C with [ $^{35}\text{S}$ ]GTP $\gamma\text{S}$  (1 nM) in the presence of vehicle, test compound (one to eight concentrations), or excess 8-OH-DPAT to define maximum agonist response. All reactions were terminated by the addition of ice cold Tris buffer followed by rapid filtration using a Tom Tech<sup>®</sup> filtration device to separate bound [ $^{35}\text{S}$ ]GTP $\gamma\text{S}$  from free compound. Agonists produce  
25 an increase in the amount of bound [ $^{35}\text{S}$ ]GTP $\gamma\text{S}$  whereas antagonists produce no increase in binding. Bound radioactivity was counted and analyzed as above.

The following assays were performed by incubating the cells with DMEM containing 25 mM HEPES, 5 mM theophylline and 10  $\mu\text{M}$  pargyline for a period of 20  
30 minutes at 37°C. Functional activity was assessed by treating the cells with forskolin (1  $\mu\text{M}$  final concentration) followed immediately by test compound (6 concentrations) for an additional 10 min at 37°C. In separate experiments, 6 concentrations of antagonist were preincubated for 20 min prior to the addition of 10 nM 8-OH-DPAT and forskolin. The reaction was terminated by removal of the media and addition of 0.5 ml

- 42 -

ice cold assay buffer. Plates were stored at -20°C prior to assessment of cAMP formation by a cAMP SPA assay (Amersham).

The compounds tested correspond to those prepared in Examples 1 to 17 above. The results of the procedure are set forth in Table 1.

TABLE 1

Example No.	5-HT <sub>1A</sub> (K <sub>i</sub> , nM)	ST(K <sub>i</sub> , nM, or %Inhib.@.1μM)	GTPγS ED <sub>50</sub> (EMax%)	cAMP ED <sub>50</sub> (EMax%)
1a	1.97	22	13.0 (80 %)	1.32 (94%)
1b	7.72	25	44.0 (60 %)	
2	1.34	0.48	41.6 (77 %)	4.28 (97%)
3a	35.0	0.97	--	4.71 (99%)
3b	47.9	49 %	(100)%	--
4	91.4	28.0	--	--
5a	1.50	0.57	2.0 (90 %)	--
5b	2.47	14	4.0 (66 %)	--
5c	4.22	18	25.8 (59%)	--
6	5.80	12.0	(100 %)	--
7	0.68	0.08	15.7 (75 %)	1.11 (90 %)
8	8.53	4.5	36 (49%)	--
9a	26.6	0.84	121 (0%)	49 (0%)
9b	29.2	4.5	--	234 (0%)
10	1.63	0.74	4.72 (80%)	--
11	1.55	3.03	2.93 (99.9%)	--
12	35.1	12.0	--	--
13a	76.9	71 %	--	--
13b	170.3	40	--	--
13c	16.8	34%	--	--
13d	9.20	26 %	--	--
14	158.4	56	--	--
15	0.54	24	3.0 (76 %)	--
16	43.9	21.0	--	--
17	14.50	16.0	236 (97.5%)	--

- 43 -

As demonstrated by the results set forth above, the compounds of the present invention are active towards 5HT1A receptors and generally elevate serotonin levels by inhibiting 5-HT transport. Accordingly, the present compounds should be useful in treating disorders related to defects in serotonin concentration.

5

The compounds of this invention may be administered orally or parenterally, neat or in combination with conventional pharmaceutical carriers. Applicable solid carriers can include one or more substances which may also act as flavoring agents, lubricants, solubilizers, suspending agents, fillers, glidants, compression aids, binders or tablet-disintegrating agents or an encapsulating material. In powders, the carrier is a finely divided solid which is in admixture with the finely divided active ingredient. In tablets, the active ingredient is mixed with a carrier having the necessary compression properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain up to 99% of the active ingredient. Any of the solid carriers known to those skilled in the art may be used with the compounds of this invention. Particularly suitable solid carriers include, for example, calcium phosphate, magnesium stearate, talc, sugars, lactose, dextrin, starch, gelatin, cellulose, methyl cellulose, sodium carboxymethyl cellulose, polyvinylpyrrolidone, low melting waxes and ion exchange resins.

20

Liquid carriers may be used in preparing solutions, suspensions, emulsions, syrups and elixirs of the compounds of this invention. The compounds of this invention can be dissolved or suspended in a pharmaceutically acceptable liquid carrier such as water, an organic solvent, a mixture of both or pharmaceutically acceptable oils or fat. The liquid carrier can contain other suitable pharmaceutical additives such as solubilizers, emulsifiers, buffers, preservatives, sweeteners, flavoring agents, suspending agents, thickening agents, colors, viscosity regulators, stabilizers or osmoregulators. Suitable examples of liquid carriers for oral and parenteral administration include water (particularly containing additives as above, e.g., cellulose derivatives, preferably sodium carboxymethyl cellulose solution), alcohols (including monohydric alcohols and polyhydric alcohols, e.g., glycols) and their derivatives and oils (e.g., fractionated coconut oil and arachis oil). For parenteral administration, the carrier can also be an oily ester such as ethyl oleate and isopropyl myristate. Sterile liquid carriers are used to prepare compositions for parenteral administration.

35

Liquid pharmaceutical compositions which are sterile solutions or suspensions can be utilized by, for example, intramuscular, intraperitoneal or subcutaneous injection. Sterile solutions can also be administered intravenously. Compositions for oral administration may be either in liquid or solid composition form.

5

Preferably, the pharmaceutical compositions containing the compounds of this invention are in unit dosage form, e.g., tablets or capsules. In such form, the compositions may be sub-divided in unit doses containing appropriate quantities of the present compounds. The unit dosage forms can be packaged compositions, for example, packeted powders, vials, ampoules, prefilled syringes or sachets containing liquids. Alternatively, the unit dosage form can be, for example, a capsule or tablet itself, or it can be the appropriate number of any such compositions in package form.

10

The therapeutically effective amount of the compounds of this invention that is administered and the dosage regimen depends on a variety of factors, including the weight, age, sex, and medical condition of the subject, the severity of the disease, the route and frequency of administration, and the specific compound employed, and thus may vary widely. However, it is believed that the pharmaceutical compositions may contain the compounds of this invention in the range of about 0.1 to about 2000 mg, preferably in the range of about 0.5 to about 500 mg and more preferably between about 1 and about 100 mg. Projected daily dosages of active compound are about 0.01 to about 100 mg/kg body weight. The daily dose can be conveniently administered two to four times per day.

20


The present invention may be embodied in other specific forms without departing from the spirit and essential attributes thereof and accordingly, reference should be made to the appended claims, rather than to the foregoing specification, as indicating the scope of the invention.

25

- 45 -

**WHAT IS CLAIMED IS:**

1. A compound of the formula:



5

wherein:

R<sub>1</sub> is hydrogen, lower alkyl or aryl;

R<sub>2</sub> is hydrogen, lower alkyl, phenyl or substituted phenyl;

10 X and Y are each, independently, hydrogen, lower alkyl, lower alkoxy, or halogen, or together combine with the carbon atoms to which they are attached to complete a cyclopentyl, cyclohexyl, phenyl, pyrrolyl, pyranal, pyridinyl, dihydrofuranyl, furanyl, dioxanyl, oxazolyl or isoxazolyl group;

Z is hydrogen, halogen or lower alkoxy; with the proviso that when X, Y or Z represent lower alkoxy, they are not present at the ortho position;

15 W is hydrogen, halogen, lower alkyl, cyano or a trifluoromethyl group; and n is 2-5; or

pharmaceutically acceptable salts thereof.

2. A compound as in claim 1, wherein:

20 R<sub>1</sub> is hydrogen, methyl or aryl;

R<sub>2</sub> is hydrogen;

X and Y are each, independently, hydrogen, halogen or lower alkoxy, or together combine with the carbon atoms to which they are attached to complete a cyclopentyl, cyclohexyl, phenyl, pyridinyl, dioxanyl, oxazolyl, furanyl or dihydrofuranyl group;

25 Z is hydrogen, halogen or lower alkoxy; with the proviso that when X, Y or Z represent lower alkoxy, they are not present at the ortho position;

W is hydrogen or halogen; and

n is 2-4; or

pharmaceutically acceptable salts thereof.

30


- 46 -

3. The compound of claim 1, which is [3-(5-Fluoro-1H-indol-3-yl)-propyl]-[2-(1H-indol-4-yloxy)-ethyl]-amine.
4. The compound of claim 1, which is [2-(1H-Indol-4-yloxy)ethyl]-[3-(1H-indol-3-yl)-propyl]-amine.
- 5 5. The compound of claim 1, which is [3-(1H-Indol-3-yl)-butyl]-[2-(1H-indol-4-yloxy)-ethyl]-amine.
- 10 6. The compound of claim 1, which is [2-(2,3-Dihydro-benzo[1,4]dioxin-5-yloxy)-ethyl]-[2-(1H-indol-3-yl)-ethyl]-amine.
7. The compound of claim 1, which is [2-(2,3-Dihydro-benzo[1,4]dioxin-5-yloxy)-ethyl]-[3-(5-fluoro-1H-indol-3-yl)-propyl]-amine.
- 15 8. The compound of claim 1, which is [2-(6-Fluorochroman-8-yloxy)-ethyl]-[2-(1H-indol-3-yl)-ethyl]-amine.
9. The compound of claim 1, which is [2-(6-Fluorochroman-8-yloxy)-ethyl]-[3-(5-fluoro-1H-indol-3-yl)-propyl]-amine.
- 20 10. The compound of claim 1, which is [2-(6-Fluorochroman-8-yloxy)-ethyl]-[2-(5-fluoro-1H-indol-3-yl)-ethyl]-amine.
- 25 11. The compound of claim 1, which is [2-(2,3-Dihydro-benzofuran-7-yloxy)-ethyl]-3-(5-fluoro-1H-indol-3-yl)-propyl]-amine.
12. The compound of claim 1, which is [2-(Benzofuran-7-yloxy)-ethyl]-[3-(5-fluoro-1H-indol-3-yl)-propyl]-amine.
- 30 13. The compound of claim 1, which is [2-(5-Fluoro-2,3-dihydro-benzofuran-7-yloxy)-ethyl]-[2-(5-fluoro-1H-indol-3-yl)-ethyl]-amine.
14. The compound of claim 1, which is [3-(5-Fluoro-1H-indol-3-yl)-propyl]-[2-(indan-4-yloxy)-ethyl]-amine.
- 35



- 47 -

15. The compound of claim 1, which is [3-(5-Fluoro-1H-indol-3-yl)-propyl]-[2-(5,6,7,8-tetrahydro-naphthalen-1-yloxy)-ethyl]-amine.
16. The compound of claim 1, which is [3-(1H-Indol-3-yl)-propyl]-[2-(naphthalen-1-yloxy)-ethyl]amine.
17. The compound of claim 1, which is [3-(1H-Indol-3-yl)-propyl]-(2-phenoxy-ethyl)-amine.
18. The compound of claim 1, which is [3-(5-Fluoro-1H-indol-3-yloxy)-propyl]-[2-(indan-5-yloxy)-ethyl]-amine.
19. The compound of claim 1, which is [3-(1H-Indol-3-yl)-propyl]-[2-(quinolin-8-yloxy)-ethyl]-amine.
20. The compound of claim 1, which is [3-(5-Fluoro-1H-indol-3-yl)-propyl]-[2-(2-methoxy-phenoxy)-2-phenyl-ethyl]-amine.
21. The compound of claim 1, which is [3-(5-Fluoro-1H-indol-3-yl)-propyl]-[2-(2-methoxy-phenoxy)-propyl]amine.
22. A pharmaceutical composition comprising a compound of the formula:



- 25 wherein:
- R<sub>1</sub> is hydrogen, lower alkyl or aryl;
- R<sub>2</sub> is hydrogen, lower alkyl, phenyl or substituted phenyl;
- X and Y are each, independently, hydrogen, lower alkyl, lower alkoxy, or halogen, or together combine with the carbon atoms to which they are attached to complete a cyclopentyl, cyclohexyl, phenyl, pyrrolyl, pyranal, pyridinyl, dihydrofuranyl, furanyl, dioxanyl, oxazolyl or isoxazolyl group;
- 30

- 48 -


Z is hydrogen, halogen or lower alkoxy; with the proviso that when X, Y or Z represent lower alkoxy, they are not present at the ortho position;

W is hydrogen, halogen, lower alkyl, cyano or a trifluoromethyl group; and

n is 2-5; or

5 pharmaceutically acceptable salts thereof.

23. A method for alleviating the symptoms of depression in a patient in need thereof comprising administering to said patient an antidepressant effective amount of a compound of the formula:



wherein:

R<sub>1</sub> is hydrogen, lower alkyl or aryl;

R<sub>2</sub> is hydrogen, lower alkyl, phenyl or substituted phenyl;

5 X and Y are each, independently, hydrogen, lower alkyl, lower alkoxy, or halogen, or together combine with the carbon atoms to which they are attached to complete a cyclopentyl, cyclohexyl, phenyl, pyrrolyl, pyran, pyridinyl, dihydrofuranyl, furanyl, dioxanyl, oxazolyl or isoxazolyl group;

Z is hydrogen, halogen or lower alkoxy; with the proviso that when X, Y or Z represent lower alkoxy, they are not present at the ortho position;

W is hydrogen, halogen, lower alkyl, cyano or a trifluoromethyl group; and

n is 2-5; or

pharmaceutically acceptable salts thereof.

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D209/14 A61K31/40

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 478 954 A (MERRELL DOW PHARMA) 8 April 1992 (1992-04-08) the whole document ---	1-22
Y	WO 96 26923 A (PF MEDICAMENT ;HALAZY SERGE (FR); PEREZ MICHEL (FR); JOHN GARETH ( ) 6 September 1996 (1996-09-06) the whole document ---	1-22
Y	SLEIGHT A J ET AL: "IDENTIFICATION OF 5-HYDROXYTRYPTAMINE1A RECEPTOR AGENTS USING A COMPOSITE PHARMACOPHORE ANALYSIS AND CHEMICAL DATABASE SCREENING" NAUNYN-SCHMIEDEBERG'S ARCHIVES OF PHARMACOLOGY, vol. 343, 1 January 1991 (1991-01-01), pages 109-116, XP000650292 the whole document --- -/--	1-22

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&amp;" document member of the same patent family

Date of the actual completion of the international search

6 September 1999

Date of mailing of the international search report

22/09/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Stellmach, J

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	NELSON, D.L.: "Structure-Activity Relationships at 5-HT <sub>1A</sub> Receptors: Binding Profiles and Intrinsic Activity" PHARMACOL. BIOCHEM. BEHAV., vol. 40, 1991, pages 1041-1051, XP002114414 the whole document	1-22
Y	GLENNON R A: "CONCEPTS FOR THE DESIGN OF 5-HT <sub>1A</sub> SEROTONIN AGONISTS AND ANTAGONISTS" DRUG DEVELOPMENT RESEARCH, vol. 26, no. 3, 1 January 1992 (1992-01-01), pages 251-274, XP000562325 ISSN: 0272-4391 the whole document	1-22
Y	US 5 436 264 A (PFISTER JURG R ET AL) 25 July 1995 (1995-07-25) cited in the application the whole document	1-22
Y	EP 0 025 727 A (SANOFI SA) 25 March 1981 (1981-03-25) the whole document	1-22
Y	US 3 371 098 A (KRALT TEUNIS ET AL) 27 February 1968 (1968-02-27) cited in the application the whole document	1-22

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0478954 A	08-04-1992	US 5189179 A	23-02-1993
		AU 641535 B	23-09-1993
		AU 8266491 A	05-03-1992
		CA 2049803 A	01-03-1992
		CN 1059717 A,B	25-03-1992
		FI 914065 A	01-03-1992
		HU 208955 B	28-02-1994
		IL 99306 A	30-03-1995
		JP 4270264 A	25-09-1992
		NO 175430 B	04-07-1994
		PT 98800 A,B	31-08-1992
		US 5559143 A	24-09-1993
		US 5387604 A	07-02-1995
WO 9626923 A	06-09-1996	FR 2731223 A	06-09-1996
		AU 4946896 A	18-09-1996
US 5436264 A	25-07-1995	NONE	
EP 0025727 A	25-03-1981	FR 2463765 A	27-02-1981
		CA 1161849 A	07-02-1984
		CA 1164464 A	27-03-1984
		JP 56099413 A	10-08-1981
		US 4404217 A	13-09-1983
		US 4510315 A	09-04-1985
US 3371098 A	27-02-1968	NONE	

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**